

String Phenomenology: past, present and future




Progress Report: The early years 1989–2012

With: Dimitri Nanopoulos, Kajia Yuan, Edi Halyo, Ben Grinstein, Keith Dienes, Claudio Coriano, Sangyeon Chang, Marco Matone, Jogesh Pati, John Ellis, Gerald Cleaver, Per Berglund, Zongan Qiu, Maxim Pospelov, Joel Walker, Richard Garavuso, Jose Isidro, Alessandro Cafarella, Sander Nooij, John Rizos, Costas Kounnas, Cristina Timirgaziu, Elisa Manno, Marco Guzzi, Ben Assel, Kyriakos Christodoulides, Mirian Tsulaia

String Phenomenology Institute, CERN, 10 July 2011

... Before



4 July 2012

After ...

DATA → STANDARD MODEL

$$\text{SU}(3) \times \text{SU}(2) \times \text{U}(1)_Y \longrightarrow \text{SU}(5) \longrightarrow \text{SO}(10)$$

$$\left[\begin{pmatrix} v \\ e \end{pmatrix} + D_L^c \right] + \left[U_L^c + \begin{pmatrix} u \\ d \end{pmatrix} + E_L^c \right] + N_L^c$$
$$\bar{5} \quad + \quad 10 \quad + \quad 1 \quad = \quad \frac{16}{16}$$

STANDARD MODEL -> UNIFICATION

ADDITIONAL EVIDENCE:

Logarithmic running, proton longevity, neutrino masses

PRIMARY GUIDES:

3 generations

SO(10) embedding

Realistic free fermionic models

'Phenomenology of the Standard Model and string unification'

- Top quark mass $\sim 175\text{--}180\text{GeV}$ PLB 274 (1992) 47
- Generation mass hierarchy NPB 407 (1993) 57
- CKM mixing NPB 416 (1994) 63 (with Halyo)
- Stringy seesaw mechanism PLB 307 (1993) 311 (with Halyo)
- Gauge coupling unification NPB 457 (1995) 409 (with Dienes)
- Proton stability NPB 428 (1994) 111
- Squark degeneracy NPB 526 (1998) 21 (with Pati)
- Minimal Superstring Standard Model PLB 455 (1999) 135
(with Cleaver & Nanopoulos)
- Moduli fixing NPB 728 (2005) 83
- Exophobia PLB 683 (2010) 306
(with Assel, Christodoulides, Kounnas & Rizos)

Other approaches

Geometrical

- Greene, Kirklin, Miron, Ross (1987)
Donagi, Ovrut, Pantev, Waldram (1999)
Blumenhagen, Moster, Reinbacher, Weigand (2006)
Heckman, Vafa (2008)
-

Orbifolds

- Ibanez, Nilles, Quevedo (1987)
Bailin, Love, Thomas (1987)
Kobayashi, Raby, Zhang (2004)
Lebedev, Nilles, Raby, Ramos-Sanchez, Ratz, Vaudrevange, Wingerter (2007)
Blaszczyk, Groot–Nibbelink, Ruehle, Trapletti, Vaudrevange (2010)
-

Other CFTs

- Gepner (1987)
Schellekens, Yankielowicz (1989)
Gato–Rivera, Schellekens (2009)
-

Orientifolds

- Cvetic, Shiu, Uranga (2001)
Ibanez, Marchesano, Rabadañ (2001)
Kiritsis, Schellekens, Tsulaia (2008)
-


Free Fermionic Construction

Left-Movers: $\psi_{1,2}^\mu, \chi_i, y_i, \omega_i$ ($i = 1, \dots, 6$)

Right-Movers

$$\bar{\phi}_{A=1,\dots,44} = \begin{cases} \bar{y}_i, \bar{\omega}_i & i = 1, \dots, 6 \\ \bar{\eta}_i & i = 1, 2, 3 \\ \bar{\psi}_{1,\dots,5} \\ \bar{\phi}_{1,\dots,8} \end{cases}$$

$$V \longrightarrow V$$



$$f \longrightarrow -e^{i\pi\alpha(f)} f$$

$$Z = \sum_{\text{all spin structures}} c(\vec{\alpha}) Z(\vec{\beta})$$

Models \longleftrightarrow Basis vectors + one-loop phases

The NAHE set:

$$1 = \{\psi^\mu, \chi^{1,\dots,6}, y^{1,\dots,6}, \omega^{1,\dots,6} \mid \bar{y}^{1,\dots,6}, \bar{\omega}^{1,\dots,6}, \bar{\eta}^{1,2,3}, \bar{\psi}^{1,\dots,5}, \bar{\phi}^{1,\dots,8}\}$$

$$S = \{\psi^\mu, \chi^{1,\dots,6}\}, \quad N = 4 \text{ Vacua}$$

$$b_1 = \{\chi^{34}, \chi^{56}, y^{34}, y^{56} | \bar{y}^{34}, \bar{y}^{56}, \bar{\eta}^1, \bar{\psi}^{1,\dots,5}\}, \quad N = 4 \rightarrow N = 2$$

$$b_2 = \{\chi^{12}, \chi^{56}, y^{12}, \omega^{56} | \bar{y}^{12}, \bar{\omega}^{56}, \bar{\eta}^2, \bar{\psi}^{1,\dots,5}\}, \quad N = 2 \rightarrow N = 1$$

$$b_3 = \{\chi^{12}, \chi^{34}, \omega^{12}, \omega^{34} | \bar{\omega}^{12}, \bar{\omega}^{34}, \bar{\eta}^3, \bar{\psi}^{1,\dots,5}\}, \quad N = 2 \rightarrow N = 1$$

$Z_2 \times Z_2$ orbifold compactification

\implies Gauge group $SO(10) \times SO(6)^{1,2,3} \times E_8$

beyond the NAHE set

Add $\{\alpha, \beta, \gamma\}$

number of generations is reduced to three

$$SO(10) \longrightarrow SU(3) \times SU(2) \times U(1)_{T_{3R}} \times U(1)_{B-L}$$

$$U(1)_Y = \frac{1}{2}(B - L) + T_{3R} \in SO(10) !$$

$$SO(6)^{1,2,3} \longrightarrow U(1)^{1,2,3} \times U(1)^{1,2,3}$$

STRING DERIVED STANDARD-LIKE MODEL (PLB278)

	ψ^μ	χ^{12}	χ^{34}	χ^{56}	$y^{3,\dots,6}$	$\bar{y}^{3,\dots,6}$	$y^{1,2}, \omega^{5,6}$	$\bar{y}^{1,2}, \bar{\omega}^{5,6}$	$\omega^{1,\dots,4}$	$\bar{\omega}^{1,\dots,4}$	$\bar{\psi}^{1,\dots,5}$	$\bar{\eta}^1$	$\bar{\eta}^2$	$\bar{\eta}^3$	$\bar{\phi}^{1,\dots,8}$
1	1	1	1	1	1,...,1	1,...,1	1,...,1	1,...,1	1,...,1	1,...,1	1	1	1	1,...,1	
S	1	1	1	1	0,...,0	0,...,0	0,...,0	0,...,0	0,...,0	0,...,0	0	0	0	0,...,0	
b_1	1	1	0	0	1,...,1	1,...,1	0,...,0	0,...,0	0,...,0	0,...,0	1,...,1	1	0	0	0,...,0
b_2	1	0	1	0	0,...,0	0,...,0	1,...,1	1,...,1	0,...,0	0,...,0	1,...,1	0	1	0	0,...,0
b_3	1	0	0	1	0,...,0	0,...,0	0,...,0	0,...,0	1,...,1	1,...,1	1,...,1	0	0	1	0,...,0

	ψ^μ	χ^{12}	χ^{34}	χ^{56}	y^3y^6	$y^4\bar{y}^4$	$y^5\bar{y}^5$	$\bar{y}^3\bar{y}^6$	$y^1\omega^5$	$y^2\bar{y}^2$	$\omega^6\bar{\omega}^6$	$\bar{y}^1\bar{\omega}^5$	$\omega^2\omega^4$	$\omega^1\bar{\omega}^1$	$\omega^3\bar{\omega}^3$	$\bar{\omega}^2\bar{\omega}^4$	$\bar{\psi}^{1,\dots,5}$	$\bar{\eta}^1$	$\bar{\eta}^2$	$\bar{\eta}^3$	
α	0	0	0	0	1	0	0	0	0	0	1	1	0	0	1	1	1 1 1 0 0	0	0	0	1 1
β	0	0	0	0	0	0	1	1	1	0	0	0	0	1	0	1	1 1 1 0 0	0	0	0	1 1
γ	0	0	0	0	0	1	0	1	0	1	0	1	1	0	0	0	$\frac{1}{2} \frac{1}{2} \frac{1}{2} \frac{1}{2} \frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{2} 0 1$

Asymmetric $BC \Rightarrow$ all untwisted moduli are projected out!

all $y_i\omega_i\bar{y}_i\bar{\omega}_i$ are disallowed

can be translated to asymmetric bosonic identifications

$$X_L + X_R \rightarrow X_L - X_R$$

moduli fixed at enhanced symmetry point

Calculation of Mass Terms

nonvanishing correlators

$$\langle V_1^f V_2^f V_3^b \cdots V_N^b \rangle$$

gauge & string invariant

“anomalous” $U(1)_A$

$$\text{Tr}Q_A \neq 0 \Rightarrow D_A = A + \sum Q_k^A |\langle \phi_k \rangle|^2$$

$$D_j = 0 = \sum Q_k^j |\langle \phi_k \rangle|^2$$


$$\langle W \rangle = \langle \frac{\partial W_N}{\partial \eta_i} \rangle = 0 \quad N = 3, \dots$$

Supersymmetric vacuum $\langle F \rangle = \langle D \rangle = 0$.

nonrenormalizable terms \rightarrow effective renormalizable operators

$$V_1^f V_2^f V_3^b \cdots V_N^b \rightarrow V_1^f V_2^f V_3^b \frac{\langle V_4^b \cdots V_N^b \rangle}{M^{N-3}}$$

Top Quark Mass Prediction



Only $\lambda_t = \langle Qt_L^c H \rangle = \sqrt{2}g$ at $N = 3$

mass of lighter quarks and leptons \rightarrow nonrenormalizable terms

$$\rightarrow \lambda_b = \lambda_\tau = 0.35g^3 \sim \frac{1}{8}\lambda_t$$

Evolve λ_t , λ_b to low energies

$$m_t = \lambda_t v_1 = \lambda_t \frac{v_0}{\sqrt{2}} \sin \beta \quad m_b = \lambda_b v_2 = \lambda_b \frac{v_0}{\sqrt{2}} \cos \beta$$

$$\text{where } v_0 = \frac{2m_W}{g_2(M_Z)} = 246 \text{ GeV} \quad \text{and} \quad v_1^2 + v_2^2 = \frac{v_0^2}{2}$$

$$m_t = \lambda_t(m_t) \frac{v_0}{\sqrt{2}} \frac{\tan \beta}{(1 + \tan^2 \beta)^{\frac{1}{2}}} \Rightarrow$$

Hierarchical top-bottom mass relation in a superstring derived standard-like model

Alon E. Faraggi

Center For Theoretical Physics, Texas A&M University, College Station, TX 77843-4242, USA
and Astroparticle Physics Group, Houston Advanced Research Center (HARC), The Woodlands, TX 77381, USA

Received 30 September 1991

I propose a mechanism in a class of superstring standard-like models which explains the mass hierarchy between the top and bottom quarks. At the trilinear level of the superpotential only the top quark gets a nonvanishing mass term while the bottom quarks and tau lepton mass terms are obtained from nonrenormalizable terms. I construct a model which realizes this mechanism. In this model the bottom quark and tau lepton Yukawa couplings are obtained from quartic order terms. I show that $\lambda_s = \lambda_b \sim |\lambda_t|$ at the unification scale. A naive estimate yields $m_t \sim 175-180$ GeV.

One of the unresolved puzzles of the standard model is the mass splitting between the top quark and the lighter quarks and leptons. Especially difficult to understand within the context of the standard model is the big splitting in the heaviest generation. Experimental limits [1] indicate the top mass to be above 80 GeV, while the bottom and tau lepton masses are found at 5 GeV and 1.78 GeV respectively. Possible extensions to the standard model are grand unified theories. Although the main prediction of GUTs, proton decay, has not yet been observed, calculations of $\sin^2 \theta_w$ and of the mass ratio m_b/m_t support their validity. Recent calculations seem to support supersymmetric GUTs versus nonsupersymmetric ones [2]. In spite of the success of SUSY GUTs in confronting LEP data [2], an understanding of the mass splitting between the top quark and the lighter quarks and leptons is still lacking. The next level in which such an understanding may be developed is in the context of superstring theory [3].

Cabibbo mixing

PLB 307 (1993) 305

Find anomaly free solution

$$M_d \sim \begin{pmatrix} \epsilon & \frac{V_2 \bar{V}_3 \Phi_{\alpha\beta}}{M^3} & 0 \\ \frac{V_2 \bar{V}_3 \Phi_{\alpha\beta} \xi_1}{M^4} & \frac{\bar{\Phi}_2^- \xi_1}{M^2} & 0 \\ 0 & 0 & \frac{\Phi_1^+ \xi_2}{M^2} \end{pmatrix} v_2,$$

$$\epsilon < 10^{-8}$$

$$\frac{V_2 \bar{V}_3 \Phi_{\alpha\beta}}{M^3} = \frac{\sqrt{5} g^6}{64 \pi^3} \approx 2 - 3 \times 10^{-4}.$$

$$\Rightarrow |V| \sim \begin{pmatrix} 0.98 & 0.2 & 0 \\ 0.2 & 0.98 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

Three generation mixing \longrightarrow

NPB 416 (1994) 63 $|J| \sim 10^{-6}$

Cleaver, Faraggi, Manno, Timirgaziu, PRD 78 (2008) 046009
Classification of F and D flat directions in FMT reduced Higgs model
No D flat direction which is F-flat up to order eight in the superpotential
no stringent flat directions to all orders

Suggesting no supersymmetric flat directions in this model (class of models)
implying no supersymmetric moduli
only remaining perturbative moduli is the dilaton

quasi-realistic model: SLM; 3 gen; $SO(10)$ embed; Higgs & $\lambda_t \sim 1$; ...
vanishing one-loop partition function, perturbatively broken SUSY
Fixed geometrical, twisted and SUSY moduli

Cleaver *etal*, $SO(10)$ and FSU5 analysis — > stringent flat directions

Classification of fermionic $Z_2 \times Z_2$ orbifolds

Basis vectors:

$$1 = \{\psi^\mu, \chi^{1,\dots,6}, y^{1,\dots,6}, \omega^{1,\dots,6} \mid \bar{y}^{1,\dots,6}, \bar{\omega}^{1,\dots,6}, \bar{\eta}^{1,2,3}, \bar{\psi}^{1,\dots,5}, \bar{\phi}^{1,\dots,8}\}$$

$$S = \{\psi^\mu, \chi^{1,\dots,6}\},$$

$$z_1 = \{\bar{\phi}^{1,\dots,4}\},$$

$$z_2 = \{\bar{\phi}^{5,\dots,8}\},$$

$$e_i = \{y^i, \omega^i | \bar{y}^i, \bar{\omega}^i\}, \quad i = 1, \dots, 6, \quad N = 4 \text{ Vacua}$$

$$b_1 = \{\chi^{34}, \chi^{56}, y^{34}, y^{56} | \bar{y}^{34}, \bar{y}^{56}, \bar{\eta}^1, \bar{\psi}^{1,\dots,5}\}, \quad N = 4 \rightarrow N = 2$$

$$b_2 = \{\chi^{12}, \chi^{56}, y^{12}, y^{56} | \bar{y}^{12}, \bar{y}^{56}, \bar{\eta}^2, \bar{\psi}^{1,\dots,5}\}, \quad N = 2 \rightarrow N = 1$$

$$\alpha = \{\bar{\psi}^{4,5}, \bar{\phi}^{1,2}\}$$

& Gauge group $SO(6) \times SO(4) \times U(1)^3 \times$ hidden

Independent phases $c_{v_j}^{[vi]} = \exp[i\pi(v_i|v_j)]$: upper block

	1	S	e_1	e_2	e_3	e_4	e_5	e_6	z_1	z_2	b_1	b_2	α
1	-1	-1	\pm										
S			-1	-1	-1	-1	-1	-1	-1	-1	1	1	-1
e_1				\pm									
e_2					\pm								
e_3						\pm							
e_4							\pm						
e_5								\pm	\pm	\pm	\pm	\pm	\pm
e_6									\pm	\pm	\pm	\pm	\pm
z_1										\pm	\pm	\pm	\pm
z_2										\pm	\pm	\pm	\pm
b_1											\pm	\pm	\pm
b_2												-1	\pm
α													

A priori 66 independent coefficients $\rightarrow 2^{66}$ distinct vacua

The twisted matter spectrum:

$$B_{\ell_3^1 \ell_4^1 \ell_5^1 \ell_6^1}^1 = S + b_1 + \ell_3^1 e_3 + \ell_4^1 e_4 + \ell_5^1 e_5 + \ell_6^1 e_6$$

$$B_{\ell_1^2 \ell_2^2 \ell_5^2 \ell_6^2}^2 = S + b_2 + \ell_1^2 e_1 + \ell_2^2 e_2 + \ell_5^2 e_5 + \ell_6^2 e_6$$

$$B_{\ell_1^3 \ell_2^3 \ell_3^3 \ell_4^3}^3 = S + b_3 + \ell_1^3 e_1 + \ell_2^3 e_2 + \ell_3^3 e_3 + \ell_4^3 e_4 \quad l_i^j = 0, 1$$

sectors $B_{pqrs}^i \rightarrow 16$ or $\overline{16}$ of $SO(10)$ with multiplicity $(1, 0, -1)$


$B_{pqrs}^i + x \rightarrow 10$ of $SO(10)$ with multiplicity $(1, 0)$

$x = \{\bar{\psi}^{1,\dots,5}, \bar{\eta}^1, \bar{\eta}^2, \bar{\eta}^3\}$ x – map \leftrightarrow spinor–vector map

Algebraic formulas for $S = \sum_{i=1}^3 S_+^{(i)} - S_-^{(i)}$ and $V = \sum_{i=1}^3 V^{(i)}$

Spinor–vector duality:

Invariance under exchange of $\#(16 + \overline{16}) < - > \#(10)$



Symmetric under exchange of rows and columns

$$E_6 : \quad 27 = 16 + 10 + 1 \quad \overline{27} = \overline{16} + 10 + 1$$

Self-dual: $\#(16 + \overline{16}) = \#(10)$ without E_6 symmetry

Spinor–Vector duality in Orbifolds:

Using the level-one $\mathrm{SO}(2n)$ characters

$$O_{2n} = \frac{1}{2} \left(\frac{\theta_3^n}{\eta^n} + \frac{\theta_4^n}{\eta^n} \right), \quad V_{2n} = \frac{1}{2} \left(\frac{\theta_3^n}{\eta^n} - \frac{\theta_4^n}{\eta^n} \right),$$

$$S_{2n} = \frac{1}{2} \left(\frac{\theta_2^n}{\eta^n} + i^{-n} \frac{\theta_1^n}{\eta^n} \right), \quad C_{2n} = \frac{1}{2} \left(\frac{\theta_2^n}{\eta^n} - i^{-n} \frac{\theta_1^n}{\eta^n} \right).$$

where

$$\theta_3 \equiv Z_f \begin{pmatrix} 0 \\ 0 \end{pmatrix} \quad \theta_4 \equiv Z_f \begin{pmatrix} 0 \\ 1 \end{pmatrix}$$

$$\theta_2 \equiv Z_f \begin{pmatrix} 1 \\ 0 \end{pmatrix} \quad \theta_1 \equiv Z_f \begin{pmatrix} 1 \\ 1 \end{pmatrix}$$

Starting from:

$$Z_+ = (V_8 - S_8) \left(\sum_{m,n} \Lambda_{m,n} \right)^{\otimes 6} (\bar{O}_{16} + \bar{S}_{16}) (\bar{O}_{16} + \bar{S}_{16}),$$

where as usual, for each circle,

$$p_{\text{L,R}}^i = \frac{m_i}{R_i} \pm \frac{n_i R_i}{\alpha'},$$

and

$$\Lambda_{m,n} = \frac{q^{\frac{\alpha'}{4}} p_{\text{L}}^2 \bar{q}^{\frac{\alpha'}{4}} p_{\text{R}}^2}{|\eta|^2}.$$

$$\text{apply } Z_2 \times Z'_2 : g \times g'$$

$$g : (-1)^{(F_1+F_2)} \delta$$

$$F_{1,2} : (\overline{O}_{16}^{1,2}, \overline{V}_{16}^{1,2}, \overline{S}_{16}^{1,2}, \overline{C}_{16}^{1,2}) \longrightarrow (\overline{O}_{16}^{1,2}, \overline{V}_{16}^{1,2}, -\overline{S}_{16}^{1,2}, -\overline{C}_{16}^{1,2})$$

$$\text{with } \delta X_9 = X_9 + \pi R_9 ,$$


$$\delta : \Lambda_{m,n}^9 \longrightarrow (-1)^m \Lambda_{m,n}^9$$

$$g' : (x_4, x_5, x_6, x_7, x_8, x_9) \longrightarrow (-x_4, -x_5, -x_6, -x_7, +x_8, +x_9)$$

Note: A single space twisting $Z'_2 \Rightarrow N = 4 \rightarrow N = 2$

$$E_7 \rightarrow SO(12) \times SU(2)$$

$$\Rightarrow \text{Analyze } Z = \left(\frac{Z_+}{Z_g \times Z_{g'}} \right) = \left[\frac{(1+g)(1+g')}{2} \right] Z_+$$



$$a = g \quad ; \quad b = g' \quad ; \quad c = gg'$$

$$\text{P.F.} = (\square + \varepsilon \tilde{\square}) = \Lambda_{m,n} \bullet (\) + \Lambda_{m,n+1/2} \bullet (\)$$

$\varepsilon = +1$ massless massive

• sector b

$$\Lambda_{p,q} \left\{ \frac{1}{2} \left(\left| \frac{2\eta}{\theta_4} \right|^4 + \left| \frac{2\eta}{\theta_3} \right|^4 \right) [P_\epsilon^+ Q_s \bar{V}_{12} \bar{C}_4 \bar{O}_{16} + P_\epsilon^- Q_s \bar{S}_{12} \bar{O}_4 \bar{O}_{16}] + \frac{1}{2} \left(\left| \frac{2\eta}{\theta_4} \right|^4 - \left| \frac{2\eta}{\theta_3} \right|^4 \right) [P_\epsilon^+ Q_s \bar{O}_{12} \bar{S}_4 \bar{O}_{16}] \right\} + \text{massive}$$

where

$$\begin{aligned}
 P_\epsilon^+ &= \left(\frac{1 + \epsilon(-1)^m}{2} \right) \Lambda_{m,n} & P_\epsilon^- &= \left(\frac{1 - \epsilon(-1)^m}{2} \right) \Lambda_m \\
 \epsilon = +1 \Rightarrow P_\epsilon^+ &= \Lambda_{2m,n} & P_\epsilon^- &= \Lambda_{2m+1,n} \\
 \epsilon = -1 \Rightarrow P_\epsilon^+ &= \Lambda_{2m+1,n} & P_\epsilon^- &= \Lambda_{2m,n}
 \end{aligned}$$

$$\text{and} \quad 12 \cdot 2 + 4 \cdot 2 = 32$$

Further :

- The spinor–vector duality in this model is realised in terms of a continuous interpolation between two discrete Wilson lines.
- The spinor–vector duality is realised in terms of a spectral flow operator that operates in the bosonic side of the heterotic string. In the case of enhanced E_6 symmetry, the spectral flow operator acts as an internal E_6 generator. When E_6 is broken the spectral flow operator induces the spinor–vector duality map.

Pati–Salam models statistics with respect to phenomenological constraints

constraint	# of models	probability	# of models
None	100000000000	1	2.25×10^{15}
+ No gauge group enhancements.	78977078333	7.90×10^{-1}	1.78×10^{15}
+ Complete families	22497003372	2.25×10^{-1}	5.07×10^{14}
+ 3 generations	298140621	2.98×10^{-3}	6.71×10^{12}
+ PS breaking Higgs	23694017	2.37×10^{-4}	5.34×10^{11}
+ SM breaking Higgs	19191088	1.92×10^{-4}	4.32×10^{11}
+ No massless exotics	121669	1.22×10^{-6}	2.74×10^9

Constraints in second column act additionally.

- A specific choice of one-loop GSO phases
- Analysis of cubic level superpotential and flat directions
- Only one Yukawa coupling at cubic level -> heavy family
- All extra colour triplets -> massive
- One light Higgs bi-doublet
- Solely MSSM below PS breaking scale

Away from the free fermionic point:

$$\begin{aligned}
Z = & \int \frac{d^2\tau}{\tau_2^2} \frac{\tau_2^{-1}}{\eta^{12}\bar{\eta}^{24}} \frac{1}{2^3} \left(\sum (-)^{a+b+ab} \vartheta \begin{bmatrix} a \\ b \end{bmatrix} \vartheta \begin{bmatrix} a+h_1 \\ b+g_1 \end{bmatrix} \vartheta \begin{bmatrix} a+h_2 \\ b+g_2 \end{bmatrix} \vartheta \begin{bmatrix} a+h_3 \\ b+g_3 \end{bmatrix} \right)_{\psi^\mu}, \\
& \times \left(\frac{1}{2} \sum_{\epsilon, \xi} \bar{\vartheta} \begin{bmatrix} \epsilon \\ \xi \end{bmatrix}^5 \bar{\vartheta} \begin{bmatrix} \epsilon+h_1 \\ \xi+g_1 \end{bmatrix} \bar{\vartheta} \begin{bmatrix} \epsilon+h_2 \\ \xi+g_2 \end{bmatrix} \bar{\vartheta} \begin{bmatrix} \epsilon+h_3 \\ \xi+g_3 \end{bmatrix} \right)_{\bar{\psi}^{1\dots 5}, \bar{\eta}^{1,2,3}} \\
& \times \left(\frac{1}{2} \sum_{H_1, G_1} \frac{1}{2} \sum_{H_2, G_2} (-)^{H_1 G_1 + H_2 G_2} \bar{\vartheta} \begin{bmatrix} \epsilon+H_1 \\ \xi+G_1 \end{bmatrix}^4 \bar{\vartheta} \begin{bmatrix} \epsilon+H_2 \\ \xi+G_2 \end{bmatrix}^4 \right)_{\bar{\phi}^{1\dots 8}} \\
& \times \left(\sum_{s_i, t_i} \Gamma_{6,6} \begin{bmatrix} h_i | s_i \\ g_i | t_i \end{bmatrix} \right)_{(y\omega\bar{y}\bar{\omega})^{1\dots 6}} \times e^{i\pi\Phi(\gamma, \delta, s_i, t_i, \epsilon, \xi, h_i, g_i, H_1, G_1, H_2, G_2)}
\end{aligned}$$

$$\Gamma_{1,1} \begin{bmatrix} h \\ g \end{bmatrix} = \frac{R}{\sqrt{\tau_2}} \sum_{\tilde{m}, n} \exp \left[-\frac{\pi R^2}{\tau_2} |(2\tilde{m} + g) + (2n + h)\tau|^2 \right]$$

Towards String Predictions

- ## 1. Low energy supersymmetry

Specific SUSY breaking patterns

→ Collider implications

- ## 2. Additional (non-GUT) gauge bosons

Proton Stability and low-scale Z'

→ Collider signatures

- ### 3. Exotic matter

In realistic string models

Unifying gauge group \Rightarrow broken by “Wilson lines”.

\Rightarrow non-GUT physical states.

⇒ Meta-stable heavy string relics

→ Dark Mater ; UHECR candidates

Conclusions

- DATA → UNIFICATION
- STRINGS → GAUGE & GRAVITY UNIFICATION
- EXPERIMENTAL PREDICTIONS ?
- FUNDAMENTAL PRINCIPLES ?
e.g. spinor–vector duality → Physics & Geometry
phase–space duality & the equivalence postulate of QM