

PAPER CODE NO.
MATH012



THE UNIVERSITY
of LIVERPOOL

SUMMER 2009 EXAMINATIONS

Bachelor of Engineering : Foundation Year

Bachelor of Science : Foundation Year

Bachelor of Science : Year 1

Bachelor of Science : Year 2

VECTORS AND INTRODUCTION TO STATISTICS

TIME ALLOWED : Three Hours

INSTRUCTIONS TO CANDIDATES

Answer ALL questions in Section A and THREE questions from Section B.
The total of the marks available on Section A is 55.

\mathbf{i} , \mathbf{j} and \mathbf{k} are unit vectors along the x , y and z axes respectively.



THE UNIVERSITY
of LIVERPOOL

SECTION A

1. Let $ABCD$ be a parallelogram. Given that $\overrightarrow{AB} = \mathbf{u}$ and $\overrightarrow{BC} = \mathbf{v}$, express each of the following in terms of \mathbf{u} and \mathbf{v} :

- (a) \overrightarrow{CD}
- (b) \overrightarrow{BD}
- (c) \overrightarrow{BP} , where P is the mid-point of \overrightarrow{CD} .

[5 marks]

2. The points P , Q and R have Cartesian coordinates $(2,3,1)$, $(3,1,-1)$ and $(4,2,3)$ respectively, where lengths are measured in metres.

Find:

- (a) the lengths of the sides of triangle PQR , correct to the nearest centimetre
- (b) $\overrightarrow{PQ} \cdot \overrightarrow{PR}$
- (c) the angles of the triangle PQR in degrees
- (d) the coordinates of the point S such that $PQSR$ is a parallelogram.

[13 marks]

3. Let $\mathbf{u} = -\mathbf{i} + 2\mathbf{j} + 3\mathbf{k}$ and $\mathbf{v} = 2\mathbf{i} - \mathbf{j} + \mathbf{k}$ where \mathbf{i} , \mathbf{j} and \mathbf{k} are mutually orthogonal unit vectors.

- (a) Find $\mathbf{u} + 2\mathbf{v}$ and $\mathbf{u} - 2\mathbf{v}$.
- (b) Find $(2\mathbf{v} - \mathbf{u}) \cdot \mathbf{v}$ and $(\mathbf{u} + 2\mathbf{v}) \cdot \mathbf{u}$.
- (c) a unit vector parallel to $(\mathbf{u} \times \mathbf{v})$
- (d) $(\mathbf{u} \times \mathbf{v}) \cdot (\mathbf{u} + 2\mathbf{v})$.

[9 marks]



THE UNIVERSITY
of LIVERPOOL

4. The points A and B have Cartesian coordinates $(2, 2, 2)$ and $(4, 4, 1)$ respectively.
- (a) Compute \overrightarrow{AB}
 - (b) Find the vector equation of the line \mathcal{L} through A and B
 - (c) the coordinates of the point at twice the distance along the line from A to B .

[5 marks]

5. Let O be a fixed origin and let \mathbf{i} , \mathbf{j} and \mathbf{k} be constant, mutually orthogonal unit vectors. A particle P moves so that its position vector \mathbf{r} with respect to O at time t is given by

$$\mathbf{r} = t^2\mathbf{i} + (t - 1)\mathbf{j} + te^t\mathbf{k}$$

where t is measured in seconds and distances are measured in metres.
Find:

- (a) the position of P at time $t = 0$ seconds
- (b) the velocity of P at time t seconds
- (c) the speed of P at $t = 2$ seconds, to the nearest cm/sec
- (d) the acceleration of P at $t = 0$.

[7 marks]



THE UNIVERSITY
of LIVERPOOL

6. An aircraft sets out from the origin O . The wind velocity relative to the ground is $\mathbf{w} = 50\mathbf{i}$ km/hr where \mathbf{i} is a unit vector pointing East.

The aircraft travels at a constant velocity $\mathbf{u} = (-150\mathbf{i} + 400\mathbf{j})$ km/hr relative to the air. Here \mathbf{j} is a unit vector pointing North.

- (a) Give an expression for the velocity \mathbf{v} of the aircraft relative to the ground.
- (b) Hence write down an expression for the position vector of the aircraft at time t hours.
- (c) Find the time in minutes at which the aircraft has flown 160 km North (relative to the ground).
- (d) Find the position vector of the point P the aircraft reaches after it has flown 160 km North (relative to the ground).

[6 marks]

7. Evaluate the determinant

$$\begin{vmatrix} 2 & 1 & x \\ 1 & 0 & 4 \\ 2 & x & -1 \end{vmatrix}.$$

Find the values of x for which the determinant is equal to 18.

[5 marks]

8. What does the conditional probability $P(X|Y)$ of events X and Y mean?

Two machines A and B make compact discs (CDs). In a given batch at the factory, 10% of the CDs are made by A and 90% by B . Also, 80% of the CDs made by A are acceptable, and 60% of the CDs made by B are acceptable. What is the probability that a given CD chosen at random from the whole batch is acceptable?

[5 marks]



THE UNIVERSITY
of LIVERPOOL

SECTION B

9. The four distinct points A, B, C and D are non-collinear and such that $\overrightarrow{AB} = \mathbf{u}$, $\overrightarrow{BC} = \mathbf{v}$ and $\overrightarrow{CD} = \mathbf{w}$. Suppose that, in terms of mutually orthogonal unit vectors \mathbf{i}, \mathbf{j} and \mathbf{k} ,

$$\mathbf{u} = 2\mathbf{i} - 3\mathbf{j} + 2\mathbf{k}, \quad \mathbf{v} = \mathbf{i} - \mathbf{j} \quad \text{and} \quad \mathbf{w} = 4\mathbf{i} - 7\mathbf{j} + 6\mathbf{k}.$$

- (a) Using the vectors \mathbf{u} and \mathbf{v} , show that a unit vector normal to the plane containing the points A, B and C is given by

$$\mathbf{n} = \frac{2\mathbf{i} + 2\mathbf{j} + \mathbf{k}}{3}.$$

- (b) Show by explicitly calculating the scalar products that

$$\mathbf{n} \cdot \mathbf{u} = 0 \quad \text{and} \quad \mathbf{n} \cdot \mathbf{v} = 0.$$

- (c) Show that A, B, C and D lie in the same plane.
(d) Suppose that A is the point $(2, 3, 2)$. What is the Cartesian equation of the plane through A, B, C and D ?
(e) Find where the straight line

$$\mathbf{r} = 2\mathbf{i} - \mathbf{j} + 4\mathbf{k} + \lambda(2\mathbf{i} - 3\mathbf{j} + 4\mathbf{k})$$

intersects the plane through A, B, C and D .

[15 marks]



THE UNIVERSITY
of LIVERPOOL

10. Suppose that the line \mathcal{L}_1 has vector equation

$$\mathbf{r} = 2\mathbf{i} + 6\mathbf{j} - 6\mathbf{k} + \lambda(\mathbf{i} - \mathbf{j} + 2\mathbf{k})$$

and that the line \mathcal{L}_2 has vector equation

$$\mathbf{r} = 3\mathbf{i} + \mathbf{j} + \mathbf{k} + \mu(2\mathbf{i} + 2\mathbf{j} - \mathbf{k}).$$

- Write down the coordinates of *any* two points on the line \mathcal{L}_1 .
- Determine two unit vectors $\hat{\mathbf{u}}_1$ and $\hat{\mathbf{u}}_2$ which are respectively parallel to the lines \mathcal{L}_1 and \mathcal{L}_2 .
- Compute the angle between the lines.
- Show that the lines intersect and find the coordinates of the point of intersection.

[15 marks]

11. The planes Π_1 , Π_2 and Π_3 have equations

$$x - 4y + z = 12, \quad 2x - 2y - z = 9, \quad \text{and} \quad x + y + 2z = 3,$$

respectively.

- Find a normal to each plane.
- Find the angle in degrees between the normals to the planes Π_1 and Π_2 .
- Find the point of intersection of Π_1 , Π_2 and Π_3 .

[15 marks]



THE UNIVERSITY
of LIVERPOOL

12. Define the mean, mode and median of a set of values.

The rainfall on 15 consecutive days is measured (in millimetres, mm)
as

5, 3, 2, 0, 5, 7, 10, 11, 1, 7, 3, 1, 2, 13, 5.

- (a) Draw a bar chart to show the number of days with rainfalls in the ranges 0-5mm, 6-10mm and 11-15mm.
- (b) What is the frequency and relative frequency of a result of 7mm?
- (c) What is the mean rainfall in millimetres?
- (d) What are the mode and median?
- (e) What is the standard deviation of the daily rainfall (to the nearest 0.01 millimetre)?

[15 marks]