

MATH423 String Theory Solutions 4

1.

$$\frac{d^2 x^\mu}{ds^2} = 0 \quad \tau = f(s). \quad (1)$$

$$\frac{dx^\mu}{ds} = \frac{dx^\mu}{d\tau} \frac{d\tau}{ds} = \frac{dx^\mu}{d\tau} f'(s) \quad (2)$$

$$\frac{d^2 x^\mu}{ds^2} = \frac{d^2 x^\mu}{d\tau^2} [f'(s)]^2 + \frac{dx^\mu}{d\tau} f''(s) \quad (3)$$

\Rightarrow equation of motion is $\frac{d^2 x^\mu}{ds^2} = 0$ if and only if $f''(s) = 0$

i.e. $f(s) = As + B$ with A, B constants.

i.e. allowed reparametrisations are a shift of origin and a rescaling.

+++++

2. We have to take the variation $x^\mu(\tau) \rightarrow x^\mu(\tau) + \delta x^\mu(\tau)$ in the action

$$S = -mc \int_{\tau_i}^{\tau_f} \sqrt{-\eta_{\mu\nu} \frac{dx^\mu}{d\tau} \frac{dx^\nu}{d\tau}} d\tau .$$

The variation gives

$$\begin{aligned} S + \delta S &= -mc \int_{\tau_i}^{\tau_f} \sqrt{-\eta_{\mu\nu} \frac{d(x^\mu + \delta x^\mu)}{d\tau} \frac{d(x^\nu + \delta x^\nu)}{d\tau}} d\tau \\ &= -mc \int_{\tau_i}^{\tau_f} \sqrt{-\eta_{\mu\nu} \left(\frac{dx^\mu}{d\tau} + \frac{d\delta x^\mu}{d\tau} \right) \left(\frac{dx^\nu}{d\tau} + \frac{d\delta x^\nu}{d\tau} \right)} d\tau \\ &= -mc \int_{\tau_i}^{\tau_f} \sqrt{-\eta_{\mu\nu} \left(\frac{dx^\mu}{d\tau} \frac{dx^\nu}{d\tau} + \frac{d\delta x^\mu}{d\tau} \frac{dx^\nu}{d\tau} + \frac{dx^\mu}{d\tau} \frac{d\delta x^\nu}{d\tau} + \frac{d\delta x^\mu}{d\tau} \frac{d\delta x^\nu}{d\tau} \right)} d\tau \\ &= -mc \int_{\tau_i}^{\tau_f} \sqrt{-\eta_{\mu\nu} \left(\frac{dx^\mu}{d\tau} \frac{dx^\nu}{d\tau} + 2 \frac{d\delta x^\mu}{d\tau} \frac{dx^\nu}{d\tau} + \frac{d\delta x^\mu}{d\tau} \frac{d\delta x^\nu}{d\tau} \right)} d\tau \end{aligned}$$

We have to Taylor expand the square root. Recall that Taylor expansion for an arbitrary function to first order is

$$f(x_0 + \delta) = f(x_0) + f'(x_0)\Delta.$$

Here,

$$f(x_0) = \sqrt{-\eta_{\mu\nu} \frac{dx^\mu}{d\tau} \frac{dx^\nu}{d\tau}} \quad \text{and} \quad \Delta = -\eta_{\mu\nu} \left(2 \frac{d\delta x^\mu}{d\tau} \frac{dx^\nu}{d\tau} + \frac{d\delta x^\mu}{d\tau} \frac{d\delta x^\nu}{d\tau} \right)$$

Hence we get (dropping terms of order δ^2)

$$\begin{aligned} S + \delta S &= -mc \int_{\tau_i}^{\tau_f} \sqrt{-\eta_{\mu\nu} \frac{dx^\mu}{d\tau} \frac{dx^\nu}{d\tau}} d\tau - mc \int_{\tau_i}^{\tau_f} \frac{1}{2} \left(\frac{-2\eta_{\mu\nu} \frac{d\delta x^\mu}{d\tau} \frac{dx^\nu}{d\tau}}{\sqrt{-\eta_{\alpha\beta} \frac{dx^\alpha}{d\tau} \frac{dx^\beta}{d\tau}}} \right) d\tau \\ \Rightarrow \quad \delta S &= -mc \int_{\tau_i}^{\tau_f} \frac{1}{2} \left(\frac{-2\eta_{\mu\nu} \frac{d\delta x^\mu}{d\tau} \frac{dx^\nu}{d\tau}}{\sqrt{-\eta_{\alpha\beta} \frac{dx^\alpha}{d\tau} \frac{dx^\beta}{d\tau}}} \right) d\tau = mc \int_{\tau_i}^{\tau_f} \frac{\frac{d\delta x^\mu}{d\tau} \frac{dx_\mu}{d\tau}}{\sqrt{-\eta_{\alpha\beta} \frac{dx^\alpha}{d\tau} \frac{dx^\beta}{d\tau}}} d\tau \end{aligned}$$

Integrating by parts ($\int V dU = \int d(VU) - U dV$) with,

$$V = \frac{\frac{dx_\mu}{d\tau}}{\sqrt{-\eta_{\alpha\beta} \frac{dx^\alpha}{d\tau} \frac{dx^\beta}{d\tau}}}, \quad U = \delta x^\mu \quad \text{we have}$$

$$\delta S = mc \int_{\tau_i}^{\tau_f} \frac{d}{d\tau} \left(\frac{\frac{dx_\mu}{d\tau}}{\sqrt{-\eta_{\alpha\beta} \frac{dx^\alpha}{d\tau} \frac{dx^\beta}{d\tau}}} \delta x^\mu \right) d\tau - \int_{\tau_i}^{\tau_f} \frac{d}{d\tau} \left[\frac{mc \frac{dx_\mu}{d\tau}}{\sqrt{-\eta_{\alpha\beta} \frac{dx^\alpha}{d\tau} \frac{dx^\beta}{d\tau}}} \right] \delta x^\mu(\tau) d\tau.$$

The first term vanishes by the conditions $\delta x^\mu(\tau_i) = \delta x^\mu(\tau_f) = 0$. Since $\delta x^\mu(\tau)$ is arbitrary in the domain of integration, The second term vanishes iff the integrand is identically zero *i.e.*

$$\frac{d}{d\tau} \left[\frac{mc \frac{dx_\mu}{d\tau}}{\sqrt{-\eta_{\alpha\beta} \frac{dx^\alpha}{d\tau} \frac{dx^\beta}{d\tau}}} \right] = 0 \quad (4)$$

This equation is in manifestly reparameterization invariant form. Indeed, the object between the brackets is clearly reparameterization invariant:

$$\frac{mc \frac{dx_\mu}{d\tau}}{\sqrt{-\eta_{\alpha\beta} \frac{dx^\alpha}{d\tau} \frac{dx^\beta}{d\tau}}} = \frac{mc \frac{dx_\mu}{d\tau'}}{\sqrt{-\eta_{\alpha\beta} \frac{dx^\alpha}{d\tau'} \frac{dx^\beta}{d\tau'}}}$$

This follows from the chain rule $\frac{dx^\mu}{d\tau} = \frac{dx^\mu}{d\tau'} \frac{d\tau'}{d\tau}$. The derivative in front of the square bracket does not spoil the reparameterization invariance since $\frac{d}{d\tau}[\dots] = \frac{d}{d\tau'}[\dots] \frac{d\tau'}{d\tau} = 0 \rightarrow \frac{d}{d\tau'}[\dots] = 0$. When we choose $\tau = s$

$$-\eta_{\mu\nu} \frac{dx^\mu}{d\tau} \frac{dx^\nu}{d\tau} = -\eta_{\mu\nu} \frac{dx^\mu}{ds} \frac{dx^\nu}{ds} = \frac{(ds)^2}{(ds)^2} = 1$$

Equation (4) then becomes

$$\frac{d}{ds} \left[mc \frac{dx_\mu}{ds} \right] = \frac{dp_\mu}{ds} = 0.$$

+++++

3. The relativistic version of Newton's second law is

$$\frac{d\vec{p}}{dt} = \frac{d}{dt} \left(\frac{m\vec{v}}{\sqrt{1-\vec{v}^2}} \right) = \vec{F}. \quad (5)$$

or in covariant form

$$\frac{dp^\mu}{ds} = \frac{d}{ds} \left(m \frac{dx^\mu}{ds} \right) = f^\mu \quad (6)$$

with

$$f^\mu = \left(\frac{\vec{v} \cdot \vec{F}, \vec{F}}{\sqrt{1-\vec{v}^2}} \right).$$

we want to show that (6) is the same as (5). Note

$$\begin{aligned} \vec{v} \cdot \frac{d\vec{p}}{dt} &= \vec{v} \cdot \frac{d}{dt} \left(\frac{m\vec{v}}{\sqrt{1-\vec{v}^2}} \right) = \frac{m\vec{v} \cdot \frac{d\vec{v}}{dt}}{(1-\vec{v}^2)^{\frac{3}{2}}} = \frac{d}{dt} \left(\frac{m}{\sqrt{1-\vec{v}^2}} \right) = \frac{dp^0}{dt} \\ &\implies \frac{dp^0}{dt} = \vec{v} \cdot \vec{F} \\ &\implies \frac{dp^0}{ds} \frac{ds}{dt} = \vec{v} \cdot \vec{F} \quad \text{or} \quad \frac{dp^0}{ds} = \frac{\vec{v} \cdot \vec{F}}{\sqrt{1-\vec{v}^2}} \end{aligned}$$

Similarly,

$$\frac{d\vec{p}}{dt} = \frac{d\vec{p}}{ds} \frac{ds}{dt} = \vec{F}$$

or

$$\frac{d\vec{p}}{ds} = \frac{\vec{F}}{\sqrt{1-\vec{v}^2}}$$

and

$$\frac{dp^\mu}{ds} = f^\mu$$

Here s is the proper time and $ds = \sqrt{1-\vec{v}^2}dt$. Note that we fixed $c = 1$.

+++++

4.

$$S = \int \frac{1}{2}mv^2 dt + \frac{q}{c} \int A_\mu(x) \frac{dx^\mu}{dt} dt \quad \text{where } A^\mu = (\Phi, \vec{A}).$$

a.

$$S = \int \frac{1}{2}mv^2 dt + \frac{q}{c} \int \vec{A} \cdot \vec{v} dt - q \int \Phi dt = \int L dt .$$

where

$$L = \frac{1}{2}mv^2 + \frac{q}{c} \vec{A} \cdot \vec{v} - q\Phi$$

b.

$$\vec{p} = \frac{\partial L}{\partial \vec{v}} = m\vec{v} + \frac{q}{c} \vec{A}$$

c.

$$H = \vec{p} \cdot \vec{v} - L = mv^2 + \frac{q}{c} \vec{A} \cdot \vec{v} - \frac{1}{2}mv^2 - \frac{q}{c} \vec{A} \cdot \vec{v} + q\Phi = \frac{1}{2}mv^2 + q\Phi$$

We have to replace \vec{v} by \vec{p} . From part (b)

$$\vec{v} = \frac{1}{m} \left(\vec{p} - \frac{q}{c} \vec{A} \right)$$

hence

$$H = \frac{1}{2m} \left(\vec{p} - \frac{q}{c} \vec{A} \right)^2 + q\Phi$$

+++++

5.

We are interested in variation of the action

$$S = -mc \int_{\mathcal{P}} ds + \frac{q}{c} I, \quad I = \int_{\mathcal{P}} d\tau A_\mu(x(\tau)) \frac{dx^\mu}{d\tau}(\tau) . \quad (7)$$

when we let $x^\mu(\tau) \rightarrow x^\mu(\tau) + \delta x^\mu(\tau)$. We note that

$$\delta A_\mu(x(\tau)) \equiv A_\mu(x(\tau) + \delta x(\tau)) - A_\mu(x(\tau)) = \frac{\partial A_\mu}{\partial x^\nu} \delta x^\nu(\tau),$$

where $\frac{\partial A_\mu}{\partial x^\nu}$ is calculated at $x = x(\tau)$. The variation of the first part was done in problem 2. The variation of the second part is obtained from

$$\begin{aligned} I + \delta I &= \int_{\mathcal{P}} d\tau A_\mu(x(\tau) + \delta x(\tau)) \frac{d}{d\tau} (x^\mu + \delta x^\mu) \\ &= \int_{\mathcal{P}} d\tau \left(A_\mu(x(\tau)) + \frac{\partial A_\mu}{\partial x^\nu} \delta x^\nu \right) \left(\frac{dx^\mu}{d\tau} + \frac{d\delta x^\mu}{d\tau} \right) \\ &= A_\mu(x(\tau)) \frac{dx^\mu}{d\tau} + \frac{\partial A_\mu}{\partial x^\nu} \delta x^\nu \frac{dx^\mu}{d\tau} + A_\mu(x(\tau)) \frac{d\delta x^\mu}{d\tau} \end{aligned}$$

where we dropped the term of order δ^2 in the last line. We therefore have

$$\delta I = \int_{\mathcal{P}} d\tau \frac{\partial A_\mu}{\partial x^\nu} \delta x^\nu \frac{dx^\mu}{d\tau} + \int_{\mathcal{P}} d\tau A_\mu(x(\tau)) \frac{d\delta x^\mu}{d\tau}.$$

We exchange $\mu \leftrightarrow \nu$ in the first term and rewrite the second using a total derivative:

$$\delta I = \int_{\mathcal{P}} d\tau \delta x^\mu \frac{\partial A_\nu}{\partial x^\mu} \frac{dx^\nu}{d\tau} + \int_{\mathcal{P}} d\tau \left[\frac{d}{d\tau} (A_\mu \delta x^\mu) - \delta x^\mu \frac{dA_\mu}{d\tau} \right].$$

We assume that δx vanishes at the ends of \mathcal{P} , so the total derivative (first term in square brackets) vanishes. Using the chain rule for the second term in square brackets we find

$$\delta I = \int_{\mathcal{P}} d\tau \delta x^\mu \left(\frac{\partial A_\nu}{\partial x^\mu} - \frac{\partial A_\mu}{\partial x^\nu} \right) \frac{dx^\nu}{d\tau} = \int_{\mathcal{P}} d\tau \delta x^\mu F_{\mu\nu} \frac{dx^\nu}{d\tau}$$

This concludes the variation of I .

The variation of the first term in (7) is obtained by varying the path $x^\mu(\tau) \rightarrow x^\mu(\tau) + \delta x^\mu(\tau)$, as was done in problem 2, or alternatively as we did in the lectures,

$$\begin{aligned} \delta S &= -mc \int \delta(ds) = mc \int \eta_{\mu\nu} \frac{d\delta x^\mu}{d\tau} \frac{dx^\nu}{ds} d\tau \\ &= mc \int_{\tau_i}^{\tau_f} d\tau \frac{d}{d\tau} \left(\eta_{\mu\nu} \delta(x^\mu(\tau)) \frac{dx^\nu}{ds} \right) - \int_{\tau_i}^{\tau_f} d\tau \delta(x^\mu(\tau)) \left(mc \eta_{\mu\nu} \frac{d}{d\tau} \left(\frac{dx^\nu}{ds} \right) \right) \end{aligned}$$

The first term is a boundary term and vanishes by imposing

$$\delta(x^\mu(\tau_i)) = \delta(x^\mu(\tau_f)) = 0.$$

$$\Rightarrow \delta S = - \int_{\tau_i}^{\tau_f} d\tau \delta(x^\mu(\tau)) \left(mc \eta_{\mu\nu} \frac{d}{d\tau} \left(\frac{dx^\nu}{ds} \right) \right)$$

The momentum four vector is given by

$$p^\nu = m u^\nu = mc \frac{dx^\nu}{ds},$$

where u^ν is the velocity four vector. Hence,

$$\delta S = - \int_{\tau_i}^{\tau_f} d\tau \delta(x^\mu(\tau)) \eta_{\mu\nu} \frac{dp^\nu}{d\tau} = - \int_{\tau_i}^{\tau_f} d\tau \delta(x^\mu(\tau)) \frac{dp_\mu}{d\tau}$$

Combining the variation of the two terms in the action for a charged particle in an electromagnetic field we get

$$\frac{dp_\mu}{d\tau} = \frac{q}{c} F_{\mu\nu} \frac{dx^\nu}{d\tau}$$