



Muyang Liu

Motivation

Visualization

Construction

2-groups

Summary

# 2-groups and T-duality in 6D Little String Theory

Muyang Liu

Uppsala University

July 7, 2022

With Michele Del Zotto, Paul-Konstantin Oehlmann – 2207-xxxx



# Outline

Muyang Liu

Motivation  
Visualization  
Construction  
2-groups  
Summary

## 1 Motivation

## 2 Visualization

## 3 Construction

## 4 2-groups

## 5 Summary



# 6D Little String Theory (LST)

Muyang Liu

Motivation

Visualization

Construction

2-groups

Summary

Has the following features

- **Decoupled from gravity**
- Compactified on **non compact** Calabi-Yau threefold  $CY_3$
- **Negative semi definite** dirac pairing  $\eta^{IJ} \geq 0$   
→ has a **zero curve**  $\Sigma_0 = \sum N_I \Sigma_I$
- String excitations have an intrinsic **string tension**  
→ key in the dynamics of the theory
- Observes **T-dualities** upon circle reduction



# 6D Little String Theory (LST)

Muyang Liu

Motivation

Visualization

Construction

2-groups

Summary

Has the following features

- **Decoupled from gravity**
- Compactified on **non compact** Calabi-Yau threefold  $CY_3$
- **Negative semi definite** dirac pairing  $\eta^{IJ} \geq 0$   
→ has a **zero curve**  $\Sigma_0 = \sum N_I \Sigma_I$
- String excitations have an intrinsic **string tension**  
→ key in the dynamics of the theory
- Observes **T-dualities** upon circle reduction
- All known LSTs can be realised within **F-theory**

[Bhardwaj, Del Zotto, Heckman, Morrison, Rudelius, Vafa '15]

⇒ Construct Heterotic ALE instantons and identify T-duals.

Detailed of geometric engineering : See Paul's talk



# Outline

Muyang Liu

Motivation

Visualization

Construction

2-groups

Summary

## 1 Motivation

## 2 Visualization

## 3 Construction

## 4 2-groups

## 5 Summary



# Geometrizing heterotic ALE instantons

Muyang Liu

Motivation

Visualization

Construction

2-groups

Summary

- The **F-theory dual** of the **Heterotic string**: an elliptic K3 with a stable degeneration limit into two  $dP_9$ s

$$\begin{array}{ccc} \mathbb{T}_f^2 & \longrightarrow & K3 \\ & \downarrow f & \rightarrow \\ & \mathbb{P}^1 & \\ & & \mathbb{T}_H^2 \\ & & \downarrow f \\ & & \mathbb{P}^1 \end{array}$$



# Geometrizing heterotic ALE instantons

Muyang Liu

Motivation

Visualization

Construction

2-groups

Summary

- The **F-theory dual** of the **Heterotic string**: an elliptic K3 with a stable degeneration limit into two  $dP_9$ s

$$\begin{array}{ccc} \mathbb{T}_f^2 & \longrightarrow & K3 \\ & \downarrow f & \rightarrow \\ & \mathbb{P}^1 & \\ & & \mathbb{T}_f^2 \longrightarrow dP_9 \vee_{\mathbb{T}_H^2} dP_9 \\ & & \downarrow f \\ & & \mathbb{P}^1 \end{array}$$

- Then, N **Heterotic instantons** are realized as a  $\mathbb{C}^2/\mathbb{Z}_N$  singularity where the two -1 curves of  $P_1$  bases intersect.



# Geometrizing heterotic ALE instantons

Muyang Liu

Motivation

Visualization

Construction

2-groups

Summary

- The **F-theory dual** of the **Heterotic string**: an elliptic K3 with a stable degeneration limit into two  $dP_9$ s

$$\begin{array}{ccc} \mathbb{T}_f^2 & \longrightarrow & K3 \\ & \downarrow f & \rightarrow \\ & \mathbb{P}^1 & \\ & & \downarrow f \\ & & \mathbb{P}^1 \end{array}$$

- Then,  $N$  **Heterotic instantons** are realized as a  $\mathbb{C}^2/\mathbb{Z}_N$  singularity where the two -1 curves of  $P_1$  bases intersect.
- The six dimensional reduction of the  $E_8 \times E_8$  Heterotic string on  $T_H^2 \rightarrow \mathbb{C}$  is equivalent to F-theory on  $K3 \rightarrow \mathbb{C}$   
 $\leftrightarrow$  construct  $CY_3$  using toric geometry [Huang Taylor'18, Anderson, Gao, Gray, Lee'16](#)



# Geometrizing heterotic ALE instantons

Muyang Liu



Motivation

Visualization

Construction

2-groups

Summary



## Hořava-Witten duality

Muyang Liu

## Motivation


## Visualization

Construction

2-groups

## Summary

- Heterotic instantons  $\leftrightarrow N$  M5 branes parallel to the two M9s
  - two copies of E8  $\leftrightarrow$  two M9s



# Heterotic strings probing ADE singularities

Muyang Liu


Motivation

## Visualization

### Construction

3 二〇一九年

- Up: M9 brane and their connecting M2 branes/strings
  - below: local F-theory curve configuration





# Outline

Muyang Liu

Motivation

Visualization

Construction

2-groups

Summary

## 1 Motivation

## 2 Visualization

## 3 Construction

## 4 2-groups

## 5 Summary



# Basic ingredients to cook a LST

Muyang Liu

Motivation

Visualization

Construction

2-groups

Summary

- Description of generalized quiver theories:

$$\mathcal{T}(\mu_1, \mathfrak{g}) \xrightarrow{\mathfrak{g}} \mathcal{T}_N(\mathfrak{g}, \mathfrak{g}) \xrightarrow{\mathfrak{g}} \mathcal{T}(\mu_2, \mathfrak{g})$$

$$\underbrace{1222\cdots221}_{N+1}$$



# Basic ingredients to cook a LST

Muyang Liu

Motivation

Visualization

Construction

2-groups

Summary

- Description of generalized quiver theories:

$$\mathcal{T}(\mu_1, \mathfrak{g}) \xrightarrow{\mathfrak{g}} \mathcal{T}_N(\mathfrak{g}, \mathfrak{g}) \xrightarrow{\mathfrak{g}} \mathcal{T}(\mu_2, \mathfrak{g})$$

$$\underbrace{1222 \cdots 221}_{N+1}$$

- $\mathcal{T}(\mu_a, \mathfrak{g})$  : M9-M5 system with  $\mathbb{C}^2/\Gamma_{\mathfrak{g}}$   
 $\leftrightarrow$  determined by choice  $\mu_a : \Gamma_{\mathfrak{g}} \rightarrow E_8$   
 $\rightarrow$  Zero form flavor symmetry.
- E.g, possible  $\mu : \mathbb{Z}_k \rightarrow E_8$  are classified by [Victor G. Kac '83](#)



# Basic ingredients to cook a LST

Muyang Liu

Motivation

Visualization

Construction

2-groups

Summary

- Description of generalized quiver theories:

$$\mathcal{T}(\mu_1, \mathfrak{g}) \xrightarrow{\mathfrak{g}} \mathcal{T}_N(\mathfrak{g}, \mathfrak{g}) \xrightarrow{\mathfrak{g}} \mathcal{T}(\mu_2, \mathfrak{g})$$

$$\underbrace{1 \ 2 \ 2 \ 2 \cdots 2 \ 2 \ 1}_{N+1}$$

- $\mathcal{T}(\mu_a, \mathfrak{g})$  : M9-M5 system with  $\mathbb{C}^2/\Gamma_{\mathfrak{g}}$   
 $\leftrightarrow$  determined by choice  $\mu_a : \Gamma_{\mathfrak{g}} \rightarrow E_8$   
 $\rightarrow$  Zero form flavor symmetry.
- E.g, possible  $\mu : \mathbb{Z}_k \rightarrow E_8$  are classified by [Victor G. Kac '83](#)
- $\mathcal{T}_N(\mathfrak{g}, \mathfrak{g})$  :  $N$  M5 branes probing a  $\mathbb{C}^2/\Gamma_{\mathfrak{g}}$  singularity;  
 $\leftrightarrow$  determined by  $[G] - [G]$  conformal matter.



Muyang Liu

Motivation


Visualization

Construction

2-groups

Summary

- M9-M5 Distance : vev of a (1,0) tensormultiplet  
↔ fractions of the intersection point on Hořava-Witten wall
- M5-M5 Distance : vev of a (1,0) tensormultiplet  
↔ fractions of M5-branes [Zotto, Heckman, Tomasiello, Vafa '14](#)





# Outline

Muyang Liu

Motivation

Visualization

Construction

2-groups

Summary

1 Motivation

2 Visualization

3 Construction

4 2-groups

5 Summary



# 2-group symmetry and T-duals

Muyang Liu

Motivation

Visualization

Construction

2-groups

Summary

- LSTs have a continuous **two-group symmetry** [Córdova,Dumitrescu, Intriligator '18]  
↔ Different form-degree symmetries mix to higher groups:

$$\left( P^{(0)} \times SU(2)_R^{(0)} \times \prod_a F_a^{(0)} \right) \times_{\widehat{\kappa}_P, \widehat{\kappa}_R, \widehat{\kappa}_{F_a}} U(1)_{LST}^{(1)}.$$



# 2-group symmetry and T-duals

Muyang Liu

Motivation

Visualization

Construction

2-groups

Summary

- LSTs have a continuous **two-group symmetry** [Córdova,Dumitrescu, Intriligator '18]  
↔ Different form-degree symmetries mix to higher groups:

$$\left( P^{(0)} \times SU(2)_R^{(0)} \times \prod_a F_a^{(0)} \right) \times_{\widehat{\kappa}_P, \widehat{\kappa}_R, \widehat{\kappa}_{F_a}} U(1)_{LST}^{(1)}.$$

- $\widehat{\kappa}_P$ ,  $\widehat{\kappa}_R$  and  $\widehat{\kappa}_{F_a}$  measure the mixing of the symmetries  
↔ must be **matched for T-duals** of LSTs.

$$\widehat{\kappa}_F = - \sum_{I=1}^{r+1} N_I \eta^{IA} \quad \widehat{\kappa}_R = \sum_{I=1}^{r+1} N_I h_{g_I}^{\vee} \quad \widehat{\kappa}_P = - \sum_{I=1}^{r+1} N_I (\eta^{II} - 2).$$



# 2-group symmetry and T-duals

Muyang Liu

Motivation  
Visualization  
Construction  
2-groups  
Summary

- LSTs have a continuous **two-group symmetry** [Córdova,Dumitrescu, Intriligator '18]  
↔ Different form-degree symmetries mix to higher groups:

$$\left( P^{(0)} \times SU(2)_R^{(0)} \times \prod_a F_a^{(0)} \right) \times_{\widehat{\kappa}_P, \widehat{\kappa}_R, \widehat{\kappa}_{F_a}} U(1)_{LST}^{(1)}.$$

- $\widehat{\kappa}_P$ ,  $\widehat{\kappa}_R$  and  $\widehat{\kappa}_{F_a}$  measure the mixing of the symmetries  
↔ must be **matched for T-duals** of LSTs.

$$\widehat{\kappa}_F = - \sum_{I=1}^{r+1} N_I \eta^{IA} \quad \widehat{\kappa}_R = \sum_{I=1}^{r+1} N_I h_{g_I}^V \quad \widehat{\kappa}_P = - \sum_{I=1}^{r+1} N_I (\eta^{II} - 2).$$

- 5d M-theory on  $CY_3$  has multiple F-theory uplifts at 6d  
↔ encoded by different elliptic fibrations in  $CY_3$ .



# Found : Multiple T-dual LSTs

Muyang Liu

Motivation

Visualization

Construction

2-groups

Summary

A concrete  $CY_3$  with **three elliptic fibrations**  $\leftrightarrow$  **T-dual triples**:

- ① An  $\mathfrak{su}_2^{M+1}$  gauge group given as the chain

$$[\mathfrak{so}_{16}^-] \underbrace{[1 \quad 2 \quad \dots \quad 2 \quad 1]}_{\times M+1} [\mathfrak{so}_{16}^+] . \quad (1)$$



# Found : Multiple T-dual LSTs

Muyang Liu

Motivation

Visualization

Construction

2-groups

Summary

A concrete  $CY_3$  with **three elliptic fibrations**  $\leftrightarrow$  **T-dual triples**:

- ➊ An  $\mathfrak{su}_2^{M+1}$  gauge group given as the chain

$$[\mathfrak{so}_{16}^-] \underbrace{\overset{\mathfrak{su}_2}{1} \overset{\mathfrak{su}_2}{2} \dots \overset{\mathfrak{su}_2}{2} \overset{\mathfrak{su}_2}{1}}_{\times M+1} [\mathfrak{so}_{16}^+]. \quad (1)$$

- ➋ A two tensor theory with

$$[\mathfrak{so}_{16}^-] \overset{\mathfrak{sp}_M}{1} \overset{\mathfrak{sp}_M}{1} [\mathfrak{so}_{16}^+]. \quad (2)$$



# Found : Multiple T-dual LSTs

Muyang Liu

Motivation

Visualization

Construction

2-groups

Summary

A concrete  $CY_3$  with **three elliptic fibrations**  $\leftrightarrow$  **T-dual triples**:

- ➊ An  $\mathfrak{su}_2^{M+1}$  gauge group given as the chain

$$[\mathfrak{so}_{16}^-] \underbrace{\overset{\mathfrak{su}_2}{1} \overset{\mathfrak{su}_2}{2} \dots \overset{\mathfrak{su}_2}{2} \overset{\mathfrak{su}_2}{1}}_{\times M+1} [\mathfrak{so}_{16}^+]. \quad (1)$$

- ➋ A two tensor theory with

$$[\mathfrak{so}_{16}^-] \overset{\mathfrak{sp}_M}{1} \overset{\mathfrak{sp}_M}{1} [\mathfrak{so}_{16}^+]. \quad (2)$$

- ➌ A single tensor theory with

$$[\mathfrak{su}_{16}] \overset{\mathfrak{su}_{2M+1}}{0}, \quad (3)$$



## More results

Muyang Liu

Motivation

## Visualization

Construction

2-groups

## Summary

$\mathfrak{g}$	$F(\mu_1, \mu_2)$	Theory description
$E_6^M$	$[E_6 \times E_6 \times SU(3)]/\mathbb{Z}_3$	$[SO(7)] \begin{smallmatrix} \mathfrak{sp}_{M-1} & \mathfrak{so}_{4M+4} & \mathfrak{sp}_{3M-3} & \mathfrak{su}_{4M} & \mathfrak{su}_{2M+6} \\ 1 & 4 & 1 & 2 & 2 \end{smallmatrix} [SU(12)]$
	$E_8 \times E_7$	$[SO(28)] \begin{smallmatrix} \mathfrak{sp}_{M+7} & \mathfrak{so}_{4M+16} & \mathfrak{sp}_{3M+1} & \mathfrak{su}_{4M+2} & \mathfrak{su}_{2M+2} \\ 1 & 4 & 1 & 2 & 2 \end{smallmatrix} [SU(2)]$
$E_7^M$	$[E_7 \times E_7 \times SU(2)]/\mathbb{Z}_2$	$[SU(16)] \begin{smallmatrix} \mathfrak{su}_{2M+10} & \mathfrak{su}_{4M+4} & \mathfrak{su}_{6M-2} & \mathfrak{sp}_{4M-4} & \mathfrak{so}_{4M+4} \\ 2 & 2 & 2 & 1 & 4 \end{smallmatrix}$
$E_8^M$	$E_7 \times E_7$	$[SO(24)] \begin{smallmatrix} \mathfrak{sp}_{M+7} & \mathfrak{so}_{4M+20} & \mathfrak{sp}_{3M+3} & \mathfrak{so}_{8M+8} & \mathfrak{sp}_{5M-3} & \mathfrak{so}_{12M-4} & \mathfrak{sp}_{4M-4} & \mathfrak{so}_{4M+4} \\ 1 & 4 & 1 & 4 & 1 & 4^* & 1 & 4 \\ [N_F=2] & & & & & & & \end{smallmatrix}$ $\frac{\mathfrak{sp}_{3M-5}}{1^*}$
$SO(4N+5)$	$E_8 \times E_8$	$[E_8] \underbrace{\dots}_{[N_F=1]} \begin{smallmatrix} \mathfrak{so}_{4N+5} & \mathfrak{sp}_{2N-2} & \mathfrak{so}_{4N+3} & \mathfrak{sp}_{2N-2-k} & \mathfrak{so}_{4N+3-2k} & \mathfrak{sp}_1 & \mathfrak{so}_9 & \mathfrak{g}_2 & \mathfrak{su}_2 \\ 1 & 4 & \dots & 1 & 4 & \dots & 1 & 4 & 1 & 3 & 2 & 2 & 1 \end{smallmatrix} [E_8]$ symmetric $k=0, \dots, 2N-3$
$SO(4N+7)$	$E_8 \times E_8$	$[E_8] \underbrace{\dots}_{[N_F=1]} \begin{smallmatrix} \mathfrak{so}_{4N+7} & \mathfrak{sp}_{2N-1} & \mathfrak{so}_{4N+5} & \mathfrak{sp}_{2N-1-k} & \mathfrak{so}_{4N+5-2k} & \mathfrak{sp}_1 & \mathfrak{so}_9 & \mathfrak{g}_2 & \mathfrak{su}_2 \\ 4 & 1 & 4 & \dots & 1 & 4 & \dots & 1 & 4 & 1 & 3 & 2 & 2 & 1 \end{smallmatrix} [E_8]$ symmetric $k=0, \dots, 2N-2$



# Outline

Muyang Liu

Motivation

Visualization

Construction

2-groups

Summary

## 1 Motivation

## 2 Visualization

## 3 Construction

## 4 2-groups

## 5 Summary



# Summary and Conclusion

Muyang Liu

Motivation

Visualization

Construction

2-groups

Summary

## Summary

- Construct various **LST** :  $F_1 \times F_2$  probing  $\mathfrak{g}$  models on non compact Calabi-Yau threefold
- **Verify** new T-duals via **2-group data matching**

## Outlook



# Summary and Conclusion

Muyang Liu

Motivation

Visualization

Construction

2-groups

Summary

## Summary

- Construct various **LST** :  $F_1 \times F_2$  probing  $\mathfrak{g}$  models on non compact Calabi-Yau threefold
- **Verify** new T-duals via **2-group data matching**

## Outlook

- Classification of heterotic strings probing ADE singularity
- Twisted compactification, genus one fibration
- More T-dual system via possible elliptic fibrations of a fixed K3